

MEDICATIONS TO STOP TAKING PRIOR TO TESTING

Please stop the following medications **5 days prior** to your allergy testing visit:

Alavert	Diphenhydramine	Pepcid
Allegra	Doxepin	Periactin
Amitriptyline	Elavil	Phenergan
Antivert	Extendryl	Polyhistine
Astelin	Famotidine	Promethazine
Astepro	Fexofenadine	Q-Dall
Atarax	Hydroxyzine	Ranitidine
Benadryl	Levocetirizine	Remeron
Cetirizine	Lodrane	Rynatan
Chlorphenamine	Loratadine	Sinequan
Cimetidine	Meclizine	Tagamet
Clarinox	Mirtazapine	Vistaril
Claritin	Nalex	Xyzal
Cyproheptadine	Olopatadine	Zantac
Desloratadine	Patanase	Zyrtec

- Do not stop any inhaled asthma medications or routine medications that treat blood pressure, heart problems, diabetes, etc.
- Please bring with you a complete list of all medications, as well as your pharmacy phone number and street address. We routinely send prescriptions electronically and will need your pharmacy information to ensure accuracy.
- Please keep in mind your initial visit will take 2-3 hours. During the testing procedure, we will be using a marking pen on your back and arms. This may cause your shirt to become stained. We suggest you bring a short sleeved t-shirt to wear after your procedure.